

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 020 527 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:

19.07.2000 Bulletin 2000/29

(51) Int. Cl.7: **C12N 15/55**

(21) Application number: 99933237.2

(86) International application number:

PCT/JP99/04167

(22) Date of filing: 03.08.1999

(87) International publication number:

WO 00/08176 (17.02.2000 Gazette 2000/07)

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

(72) Inventors:

- HAMADA, Kazuyuki,
Japan Tobacco Inc.
Shizuoka 438-802 (JP)
- NAKAKIDO, Fumio,
Japan Tobacco Inc.
Shizuoka 438-0802 (JP)

(30) Priority: 04.08.1998 JP 22006098

(71) Applicant: Japan Tobacco Inc.

Tokyo 105-8422 (JP)

(74) Representative:

Reinhard - Skuhra - Weise & Partner
Postfach 44 01 51
80750 München (DE)

(54) **MUTATED BARNASE GENE AND PLANT TRANSFORMED BY THE SAME**

(57) A male sterile plant which is free from any undesirable character is constructed by expressing a barnase gene alone specifically in the anther. The barnase gene, which has been depressed by mutagenesis while sustaining its activity, is transferred into a plant and then expressed specifically in the anther to thereby efficiently give the aimed male sterile transformant.

EP 1 020 527 A1

Description

[0001] The present application claims priority from Japanese Patent Application No.: Hei 10-220060, the disclosure of which is incorporated herein by reference.

FIELD OF THE INVENTION

[0002] This invention relates to a mutant barnase gene which makes it possible to efficiently yield a male sterile transgenic plant when expressed in a specific site of a plant, in particular, anther-specifically. The present invention further relates to a recombinant vector capable of expressing the mutant barnase gene of the invention in host cells, a plant transformed by this vector and a method for constructing a transgenic plant.

PRIOR ART

[0003] Barnase is an RNase originating in *Bacillus amyloliquefaciens* (S. Nishimura and M. Nomura, Biochem. Biophys. Acta 30, 430-431:1958; R.W. Hartley, J. Mol. Biol., 202, 913-915:1988). This enzyme has 110 amino acid residues and hydrolyzes RNA. When expressed in cells, this enzyme inhibits the functions of the cells as a result of its potent RNase activity and thus causes cell death in many cases. By using this characteristic, it is therefore expected that the function of the specific site can be selectively controlled by expressing the barnase gene in a specific site of a plant.

[0004] PCT International Publication WO89/10396 discloses a technique whereby a male sterile plant is obtained by constructing a male sterility gene by ligating the above-described barnase gene to the downstream of an anther tapetal cell-specific expression promoter and introducing the thus obtained gene into a plant. Male sterilization techniques are of great value in efficiently developing an F1 hybrid variety.

[0005] When the barnase gene is employed as a male sterility gene, however, it is frequently observed that resulting male sterile transgenic plants exhibit unfavorable characteristics. PCT International Publication WO96/26283 refers to this problem in rice. It is also reported that similar phenomena are observed not only in rice but in lettuce (Scientia Horticulturae 55, 125-139:1993; Arlette Reymaerts, Hilde Van de Wiele, Greta De Sutter, Jan Janssens: Engineered genes for fertility control and their application in hybrid seed production). According to this report, a plant with depressed activity was constructed by introducing a male sterility gene comprising a tobacco anther-specific promoter (TA29) and a barnase gene into lettuce.

[0006] Although reasons for these phenomena have not been accurately clarified so far, it is assumed that the so-called "position effect" of the site in which the gene is transferred may account for the mechanism (PCT International Publication WO96/26283). More specifically, a desired male sterile plant should be constructed if the male sterility gene is expressed exclusively in the target site (i.e., anthers). However, there is a possibility that the barnase might be expressed also in tissues other than anthers although in a very small dose under the action of expression regulators (for example, an endogenous enhancer) existing in the vicinity of the gene transfer site. In such a case, the unfavorable characteristics as described above can appear because barnase has a strong enzymatic activity.

[0007] To overcome this problem, the method disclosed in PCT International Publication WO96/26283 exploits the character of cauliflower mosaic virus 35S promoter (hereinafter referred to as CaMV35S promoter) of being expressed potently in tissues other than anthers. Namely, barstar, i.e., an inhibitory protein against barnase is employed therein. The barstar gene ligated to a CaMV35S promoter is transferred into a plant simultaneously with a barnase gene and then the barstar gene is constitutively expressed in tissues other than the anthers, thereby eliminating the effects of the barnase in tissues other than the anthers. However, it is necessary in this method to transfer not only a barnase gene but also a barstar gene, and hence, the problem is that the application of this technique to the breeding of F1 varieties of rice or corn seeds may give rise to a gene silencing. "Gene silencing" is a phenomenon wherein the expression of a gene is inhibited when plural copies of the foreign gene are introduced. It is known that this problem frequently occurs when the expression of a foreign gene is effected by a 35S promoter, though the detailed mechanism thereof has not been clarified yet (R.B. Flavell, Proc. Natl. Acad. Sci. USA 91, 3490-3496:1994; J. Finnegan, Bio. Technology 12, 883-888:1994; M.A. Matzke and A.J.M. Matzke, Plant Physiol., 107, 679-685:1995). Since, in rice and corn, a barstar gene is used as a "fertility-restoring gene" in the pollen parent (father) so as to allow the pollen in the F1 generation to restore the fertility (C. Mariani, et al., Nature 357, 384-387:1992), if an MS plant (mother) is constructed by the method with the use of a barstar gene as reported in WO96/26283, the F1 plant has plural copies of the barstar gene. In such a case, there is a possibility that the expression of the gene may be inhibited due to the gene silencing.

SUMMARY OF THE INVENTION

[0008] The present invention provides a method for constructing a male sterile plant by using a barnase gene with-

out resort to a barstar gene.

[0009] The present invention further provides a mutant barnase gene to be used in the above method and a process for producing the same.

5 DETAILED DESCRIPTION OF THE INVENTION

[0010] In the present invention, the DNA sequence of barnase gene (R.W. Hartley, J. Mol. Biol. 202, 913-915:1988) is mutated at least in part and then the thus obtained mutant barnase gene is anther-specifically expressed in a plant so as to make the plant substantially male sterile without any substantially disadvantageous effect on the tissues other than anthers.

[0011] The mutation can be performed by a known method such as site-specific mutagenesis, deletion of a fragment by using restriction enzymes or the low fidelity PCR method. Among all, it is preferred to use the low fidelity PCR method. This method is described in detail in D. Leung, E. Chen and D. Goedda, Technique 1, 11-15:1989; Y.Z. Xiaoping and R.H. Ebricht, Nucleic Acid Res. 19, 6052:1991; G.C. Rice et al., Proc. Natl. Acad. Sci. USA 89, 5467-5471: 15 1992 and the contents of these documents are incorporated herein by reference. By using this technique, PCR is carried out under such conditions as to induce some errors during the amplification reaction. Thus, random mutations can be introduced into the target DNA fragment.

[0012] Primers to be used in the low fidelity PCR method in the present invention are selected as in the conventional PCR method. It is preferred that each of these primers has a similar number of nucleotides as in the conventional PCR method.

[0013] The present inventors performed PCR by using a DNA containing the sequence represented by SEQ ID NO:1 as a template with the use of a combination of primer 1 (5'-CGTTCGGGCTC GATGGTACCG GTTATCAACA CGTTTGA-3'; SEQ ID NO: 6) and primer 2 (5'-CCTCTAGATT ATCTGATTTT TGTAAGGTC TGATAATG-3'; SEQ ID NO:7) under such conditions as to induce errors. Thus, a mutant barnase gene having the DNA sequence represented by SEQ ID NO:3 was isolated. The sequence represented by SEQ ID NO:1 employed herein is the sequence of the coding region of the barnase gene contained in the known plasmid pVE108 (WO92/ 13956). This sequence corresponds to the wild type barnase gene from which the unnecessary moiety corresponding to the secretion signal in the N-terminal side has been deleted.

[0014] It is also possible to acquire different mutant barnase genes by a similar method.

[0015] The PCR amplification product is cloned in a host (for example, *Escherichia coli*) and clones containing a mutant barnase gene is isolated. In this cloning, clones having a mutant barnase gene may be screened by assaying the RNase activity expressed by the gene. However, it is advantageous to screen the clone by the method consisting of the following two steps by taking advantage of the fact that the RNase activity of barnase affects the growth of *E. coli*.

[0016] In the first step, a plasmid is prepared by using the mutant barnase gene obtained above and then *E. coli* is transformed thereby. The growth of the *E. coli* transformant thus obtained will be inhibited by the activity of the mutant barnase. Based on this fact, a colony growing slowly (i.e., a small colony), compared with *E. coli* having a control vector free from any barnase gene, is selected. It is expected that the thus selected *E. coli* strain will contain the mutant barnase gene. To confirm that the inhibition of the growth of *E. coli* is due to the expression of the mutant barnase gene, the second step is then carried out.

[0017] In the second step, a barstar gene is employed. As described above, barstar is an inhibitory protein against barnase. In *E. coli* which expresses barstar, the enzymatic activity of barnase is inhibited and thus the degradation of mRNA, which otherwise takes place in the presence of barnase, can be inhibited. Therefore, *E. coli* which expresses a barstar gene is transformed with a barnase gene, the growth will not be inhibited. Accordingly, it is expected that the thus obtained *E. coli* transformant will produce little difference in growth rate from the transformant constructed by the control plasmid which is free from any barnase gene and, therefore, these transformants will produce little difference in colony size too. Based on this principle, a plasmid is prepared from the *E. coli* strain selected in the first step. The plasmid is then used to transform another *E. coli* strain, in which the barstar gene is constitutively expressed. Thus, mutant barnase gene-containing colonies having almost the same size as the one of the *E. coli* transformed by the control plasmid are selected. The *E. coli* with the constitutive expression of the barstar gene to be used herein can be prepared by, for example, the method as will be described in Example 1 hereinafter.

[0018] The DNA sequence of the thus obtained mutant barnase gene may be analyzed by a conventional method, if necessary, to thereby examine the mutation in detail.

[0019] A preferred example of the mutant barnase gene of the present invention is one which has the DNA sequence represented by SEQ ID NO:3. Compared with the nucleotide sequence of SEQ ID NO:1 coding for the wild type activity, the nucleotide sequence encoding this gene has an insertion of "T" at the 15-position from A in the initiation codon ATG and a deletion of "A" at the 333-position. According to the normal translation manner, translation will be initiated from ATG at the 1-position of this DNA sequence represented by SEQ ID NO:3, and terminated at the ninth codon which has been converted to a termination codon as a result of the insertion of the "T" at the 15-position. It is

unlikely that the thus formed oligopeptide consisting of 8 amino acids has the barnase activity. Further, there is no other ATG or GTG codon from which the translation can be initiated in the correct frame in the vicinity thereof. However, the fact that the growth rate of *E. coli* was restored by the barnase protein strongly suggests that the translation of the barnase protein from the gene of SEQ ID NO:3 proceeded in the correct frame. This is seemingly due to a phenomenon called "frame shift re-coding" whereby the ribosome controls the translation while automatically shifting the reading frame during the course of translation into the protein. This phenomenon, which has been reported in several genes of viruses, *E. coli* and animals, generally depends exclusively on an individual nucleotide sequence and neither any special protein nor translation device is required. The frame shifting efficiency during the translation varies from gene to gene, i.e., ranging from several % to 50%. In the case of the mutant barnase gene, it is considered that the efficiency of the shifting of the reading frame during the translation is not high and thus the translation product in the correct frame is formed in a smaller amount, thereby lowering the activity of the gene.

[0020] Moreover, there is a high possibility that other mutant barnase genes of which the translation efficiency is lowered due to the shifting of the reading frame, as in the case of the DNA sequence represented by SEQ ID NO:3, or some other reasons can be obtained by mutating the barnase gene by, for example, the low fidelity PCR method. Furthermore, there is a high possibility that a gene having a DNA sequence derived from the DNA sequence represented by SEQ ID NO:3 by substitution, deletion, insertion or addition of one or more nucleotides will be a mutant barnase gene which provides a lowered translation efficiency due to the shifting of the reading frame as a result of the insertion of T at the 15-position from A in the initiation codon ATG and/or the deletion of A at the 333-position, as in the DNA sequence represented by SEQ ID NO:3. It is considered that each of these genes, likewise the gene of SEQ ID NO:3, is capable of making a plant substantially male sterile when it is anther-specifically expressed in the plant. Accordingly, these genes are also included in the scope of the present invention similar to the gene of SEQ ID NO:3.

[0021] The plant which is to be male sterilized by the mutant barnase gene is not particularly limited, so long as the gene can be transferred into the plant to provide a male sterilized plant. Examples of the plants include rice, corn, tobacco, lettuce and Brassica. Among all, rice and corn are preferred.

[0022] In order to male sterilize a plant, the mutant barnase gene is expressed specifically in the anthers of the plant to inhibit the anther functions by the RNase activity of barnase. Such an anther-specific expression can be achieved by using the method described in WO92/13957. In brief, the mutant barnase gene is ligated to the downstream of an anther-specific promoter and then incorporated into plant cells with the use of an expression vector.

[0023] The incorporation may be performed by using the Agrobacterium method, the electroporation method, the particle gun method, etc.

[0024] From the transformed plant cells, a complete plant can be formed by the regeneration from a callus of the transformed plant cells in accordance with, for example, the method reported in Y. Hiei et al., Plant J. 6, 271-282:1994.

[0025] The male sterility of the thus constructed plant can be confined by the inability of the plant to fertilize unless it is pollinated from another fertilizable plant.

Example 1: Preparation of mutant barnase gene

Low fidelity PCR

[0026] Primer 1 (CGTTCGGCTC GATGGTACCG GTTATCAACA CGTTTGA: SEQ ID NO: 6) and Primer 2 (CCTCTAGATT ATCTGATTTT TGTAAGGTC TGATAATG: SEQ ID NO:7) were synthesized by a DNA synthesizer (manufactured by Applied Biosystems) in accordance with the method described in S.L. Beaucage et al., Tetrahedron Lett., 22, 1859-1862:1982. Known plasmid pVE108 (WO92/13956) was used as a template and PCR was carried out with the use of the combination of Primers 1 and 2. SEQ ID NO:1 shows the sequence of the coding region of the barnase gene which is contained in pVE108. The reaction was performed for 50 cycles with each cycle consisting of 1 minute at 94°C, 1 minute at 57°C and 1 minute at 72°C in 10 mM Tris-HCl (pH 9.5), 50 mM KCl, 2 mM MgCl₂, and 1 mM each of dNTP, 10 ng of the template DNA and 0.5 U of Taq DNA polymerase.

[0027] After the completion of the reaction, the amplification product was separated by electrophoreses on agarose gel (2% SeaKem GTG agarose, 1xTAE) and purified by the DEAE-cellulose method (M. Muramatsu, Labo-manual Idenshikogaku (Gene Engineering Labo-Manual), Maruzen, pp.111:1988). By using the purified product thus obtained as a template, PCR was performed again under the conditions as defined above.

Ligation of mutated barnase gene fragment into plasmid vector

[0028] The reaction product from the above step was digested with SacI and XbaI in a conventional manner and then purified by agarose gel electrophoresis to give an "insertion fragment". A plasmid to be used for transferring this insertion fragment into *E. coli* can be appropriately selected. For example, use may be made of plasmid pHM1 therefor. The plasmid pHM1 was constructed as follows. Plasmid pBR322 was cleaved at the EcoRI site and blunt-ended by

using T4 DNA polymerase (manufactured by Takara Shuzo). Separately, lacZ expression cassette (322 bp) was excised from pUC18 with PvuII and the ends thereof were blunted. The cassette was integrated into the restriction site of the blunt-ended plasmid pBR 322 to give plasmid pHM1.

5 [0029] Plasmid pHM1 was digested with SacI and XbaI followed by dephosphorylation with calf intestine alkaline phosphatase (manufactured by Takara Shuzo) to give a restriction enzyme-treated plasmid fragment. Into this restriction enzyme-treated plasmid pHM1, the above "insertion fragment" containing a barnase gene was ligated by using Takara Ligation Kit ver.1 (manufactured by Takara Shuzo).

Introduction into *E. coli* and selection of barnase-active clone

10 [0030] The *E. coli* having the barnase gene introduced thereto may be selected by, for example, the following method. Since mRNA is degraded in cells by the barnase activity, the synthesis of protein is suppressed and, in its turn, the growth of *E. coli* is inhibited. Thus, the *E. coli* transformed by the barnase gene can be selected by taking advantage of the fact that the desired transformant colony has a smaller size than the *E. coli* colony transformed by the control plasmid free from the mutant barnase gene. When the wild type barnase or a mutant barnase still maintaining the activ-
15 ity comparable thereto is integrated into pHM1, *E. coli* cannot form any colony. Thus, clones having a sufficiently weakened barnase activity can be exclusively selected.

[0031] Thus, the plasmid having the mutant barnase gene ligated thereto was precipitated from ethanol and then introduced into *E. coli* LE392 strain by the electroporation method with the use of GenePulser (BioRad) in accordance with the manufacture's instruction. Also, the control plasmid free from mutant barnase was introduced into the *E. coli* LE392 strain in the same manner.

20 [0032] Next, these *E. coli* transformants were plated onto an LB agar medium containing tetracycline (25 µg/ml) and incubated at room temperature (25°C) for 72 hours. Then colonies of a smaller size than the colony of *E. coli* having the control plasmid pHM1 introduced thereto were screened as colonies of the *E. coli* transformants having the mutant barnase transferred thereto (Table 1, A).

Selection of mutant barnase clone by barstar gene

[0033] As described above, barstar is an enzyme acting as an antagonist to barnase. In *E. coli* wherein barstar is expressed, the enzymatic activity of barnase is inhibited and thus the degradation of mRNA, which otherwise takes place in the presence of barnase, can be inhibited. Therefore, if *E. coli* expressing barstar is transformed by a plasmid containing a barnase gene, the growth thereof will not be inhibited. Accordingly, it is expected that the thus obtained *E. coli* transformant will show little difference in growth rate from the transformant constructed by using the control plasmid free from a barnase gene and, therefore, these transformants will show little difference in colony size too.

35 [0034] An *E. coli* transformant which expressed barstar was constructed in the following manner. A barstar gene described in R.W. Hartley, J. Mol. Biol. 202, 913-915:1998 was excised with HindIII and XbaI and ligated in frame to a tac promoter (de Boer et al., Proc. Natl. Acad. Sci. USA 80, 21-25:1983). The gene, together with a chloramphenicol tolerance gene (N.K. Alton and D. Vapnek, Nature 282, 864-869:1979), was ligated into the defective transposon which lacked the transferase in a vector described in Herrero et al., J. Bacteriol. 172, 6557-6567:1990. Subsequently, the thus
40 obtained plasmid was introduced into *E. coli* MC1061 strain thereby to transfer the transposon, which contained the barstar gene cassette, into the chromosome of the *E. coli*. It was confirmed based on the chloramphenicol tolerance that this *E. coli* was able to maintain the barstar gene cassette in a stable state. Since the *E. coli* MC1061 strain lacked the lacI gene, the tac promoter was continuously active and thus the barstar gene was constitutively expressed.

[0035] The plasmid containing the mutant barnase gene as the insertion fragment and the control plasmid free from the same were each introduced into the above *E. coli* transformant by the electroporation method. Then the *E. coli* transformants were plated onto an LB agar medium containing IPTG (1mM) and tetracycline (25 µg/ml) and incubated at 25°C for 72 hours. Among the colonies of the *E. coli* transformed by the mutant barnase gene, a colony being comparable in size to the colony of the *E. coli* having the control plasmid pHM1, which had been simultaneously plate-cultured, was screened and named "#4-31" (Table 1, B).

Table t

Screened clone and colony size (mm)		
	Screened clone	pHM1 (control)
A: <i>E. coli</i> LE392* ¹	1.77 ± 0.33	2.65 ± 0.47
B: <i>E. coli</i> with barstar gene expression* ²	1.39 ± 0.10	1.55 ± 0.08

*1: Colony size after incubation on the LB agar plate containing 25 µg/mL of tetracycline at 28°C for 48 hours.

*2: Colony size after incubation on the LB agar plate containing 25 µg/mL of tetracycline at 28°C for 24 hours.

Example 2: Determination of nucleotide sequence of mutant barnase gene

[0036] From the *E. coli* screened in Example t, a cloned mutant barnase gene appropriate for the object of the present invention was prepared. First, the clone #4-31 screened in Example t was propagated. Next, the mutant barnase gene fragment in the clone #4-31 was excised with KpnI and XbaI and ligated into the KpnI, XbaI-site of pUC119 having been cleaved similarly with KpnI and XbaI. After amplifying the obtained plasmid by using *E. coli*, a reaction was performed by the cycle sequence method with the use of Taq polymerase (Taq Dye Terminator Cycle Sequencing Kit, manufactured by Applied Biosystems Inc.) in accordance with the manufacturer's protocol. Subsequently, the sequence was analyzed with a DNA Sequencer (Model 373A, manufactured by Applied Biosystems Inc.), thereby yielding the sequence represented by SEQ ID NO:3. Compared with the DNA sequence of the wild type barnase gene, this sequence had an insertion of T at the 15-position from A of the initiation codon ATG and a deletion of A at the 333-position.

Example 3: Construction of male sterile rice by using attenuated mutant barnase gene

[0037] The attenuated mutant barnase gene integrated into pUC119 as described above was cleaved with restriction enzymes XbaI and KpnI and thus the plasmid vector pTS431 represented by SEQ ID NO:5 was constructed. Although this plasmid pTS431 differs from the known plasmid pVE108 (PCT International Publication WO92/13956) in the following points, it is substantially equivalent thereto except for the moieties of the anther-specific promoter and the mutant barnase.

(1) In the plasmid vector pTS431 according to the present invention, the barnase gene (SEQ ID NO:1) in the known plasmid pVE108 has been converted into the mutant barnase gene (SEQ ID NO:3).

(2) A tobacco-origin anther specific promoter is used in the known plasmid pVE108, while the anther specific promoter of rice E1 gene (PCT International Publication WO92/13956) is used in the plasmid vector pTS431 according to the present invention.

(3) In the present invention, a 35S3 promoter (EP 0344029) of 1376 bp, which is not used in the plasmid pVE108, is employed upstream of the barnase gene.

(4) In the present invention, a sequence originating in the downstream of Agrobacterium T-DNA gene 7 excised from pJD884 (PCT International Publication WO93/09218) is employed whereas such a sequence is not used in the plasmid pVE108.

(5) In the present invention, the region corresponding to lacZ has been deleted from the pUC19-origin moiety. In addition a plasmid (pTS172) having the wild type barnase gene integrated therein is represented by SEQ ID NO:4.

[0038] The fragment of about 4.5 kbp was excised with a restriction enzyme EcoRI respectively from pTS431 (a plasmid having the mutant barnase gene transferred therein; SEQ ID NO:5) and pTS172 (a plasmid having the barnase gene transferred therein; SEQ ID NO:4) and inserted into the EcoRI-site of the intermediate vector pSB1 (T. Komari et al., Plant J. 10(1), 165-174:1996), and further, the T-DNA region thereof was integrated into the acceptor vector pSB1 (T. Komari et al., Plant J. 10(1), 165-174:1996) by way of homologous recombination. By using *Agrobacterium tumefaciens* LBA4404 having the obtained recombinant plasmids (pSB1431 and pSB1172 respectively), rice (variety: Asanohikari) was transformed. While the transformation was performed basically in accordance with the method of Hiei et al. (Plant J. 6(2), 271-282:1994), phosphinothricine (concentration: 10 mg/L) was employed for screening transformants since the thus constructed male sterility genes contained bar gene encoding phosphinothricine acetyl transferase as a selective marker. Phosphinothricine facilitates selection of calluses having the gene transferred therein.

[0039] A comparison of the rice transformant obtained with the use of the wild type barnase gene with the one obtained with the use of the mutant barnase gene indicates that the transfer of the mutant barnase gene gave rise to a remarkable improvement in the transformation efficiency and an increase in the rate of morphologically normal male sterility transformants, as shown in Table 2.

Table 2

Transformation efficiency				
	No. of infected calluses	No. of regenerated calluses	No. of PCR-positive lines* ¹	No. of morphologically normal male sterile lines
pSB1172 (control)	2838	83	52/83	9/52(17.3%)
pSB1431 (present invention)	787	69	43/45* ²	27/28* ³ (96.4%)

*1: detection of barnase gene fraction by PCR.

*2: examined only 45 lines among 69.

*3: examined only 28 lines among 43.

Effects of the Invention

[0040] In the present invention, a barnase gene having a weakened effect is constructed via mutation. A male sterility gene comprising the mutant barnase gene can be introduced into a plant to successfully produce a male sterile plant free from any unfavorable characteristic at a high efficiency with the use of a single gene without resort to the use of a barstar gene.

SEQUENCE LISTING

5 <110> Japan Tobacco Inc.

<120> MUTANT BARNASE GENE AND TRANSGENIC PLANT TRANSFORMED BY SAID GENE

10 <130> 980687

15 <160> 7

<210> 1

20 <211> 343

<212> DNA

<213> Bacillus amyloliquefaciens

25 <220>

<221> NAME/KEY: CDS

30 <222> 1..336

<400> 1

35 atg gta ccg gtt atc aac acg ttt gac ggg gtt gcg gat tat ctt cag 48
Met Val Pro Val Ile Asn Thr Phe Asp Gly Val Ala Asp Tyr Leu Gln
1 5 10 15

40 aca tat cat aag cta cct gat aat tac att aca aaa tca gaa gca caa 96
Thr Tyr His Lys Leu Pro Asp Asn Tyr Ile Thr Lys Ser Glu Ala Gln
20 25 30

45 gcc ctc ggc tgg gtg gca tca aaa ggg aac ctt gca gac gtc gct ccg 144
Ala Leu Gly Trp Val Ala Ser Lys Gly Asn Leu Ala Asp Val Ala Pro
35 40 45

50 ggg aaa agc atc ggc gga gac atc ttc tca aac agg gaa ggc aaa ctc 192
Gly Lys Ser Ile Gly Gly Asp Ile Phe Ser Asn Arg Glu Gly Lys Leu

55

EP 1 020 527 A1

50 55 60

5 cgc ggc aaa agc gga cga aca tgg cgt gaa gcg gat att aac tat aca 240
Pro Gly Lys Ser Gly Arg Thr Trp Arg Glu Ala Asp Ile Asn Tyr Thr

65 70 75 80

10 tca ggc ttc aga aat tca gac cgg att ctt tac tca agc gac tgg ctg 288
Ser Gly Phe Arg Asn Ser Asp Arg Ile Leu Tyr Ser Ser Asp Trp Leu

85 90 95

15 att tac aaa aca acg gac cat tat cag acc ttt aca aaa atc aga taa 336
Ile Tyr Lys Thr Thr Asp His Tyr Gln Thr Phe Thr Lys Ile Arg

100 105 110

20 ggtaacc 343

<210> 2

25 <211> 112

<212> PRT

30 <213> Bacillus amyloliquefaciens

<400> 2

35 Met Val Pro Val Ile Asn Thr Phe Asp Gly Val Ala Asp Tyr Leu Gln

1 5 10 15

Thr Tyr His Lys Leu Pro Asp Asn Tyr Ile Thr Lys Ser Glu Ala Gln

40 20 25 30

Ala Leu Gly Trp Val Ala Ser Lys Gly Asn Leu Ala Asp Val Ala Pro

35 40 45

45 Gly Lys Ser Ile Gly Gly Asp Ile Phe Ser Asn Arg Glu Gly Lys Leu

50 55 60

Pro Gly Lys Ser Gly Arg Thr Trp Arg Glu Ala Asp Ile Asn Tyr Thr

50 65 70 75 80

Ser Gly Phe Arg Asn Ser Asp Arg Ile Leu Tyr Ser Ser Asp Trp Leu

55

EP 1 020 527 A1

	85	90	95
5	Ile Tyr Lys Thr Thr Asp His Tyr Gln Thr Phe Thr Lys Ile Arg		
	100	105	110
10	<210> 3		
	<211> 342		
	<212> DNA		
15	<213>		
	<400> 3		
20	atggtaccgg ttattcaaca cgtttgacgg ggttgcggat tatcttcaga catatcataa	60	
	gtacctgat aattacatta caaatcaga agcacaagcc ctccgctggg tggcatcaaa	120	
	agggaaacctt gcagacgtcg ctccggggaa aagcatcggc ggagacatct tctcaaacag	180	
25	ggaaggcaaa ctcccgggca aaagcggacg aacatggcgt gaagcggata ttaactatac	240	
	atcaggettc agaaattcag accggattct ttactcaagc gactggctga ttacaaaac	300	
30	aacggaccat tatcagacct ttacaaaat cagtaatcta ga	342	
	<210> 4		
35	<211> 6548		
	<212> DNA		
	<213> <u>Escherichia coli</u> LE392		
40	<220>		
	<223> Clone: pTS172		
45	<400> 4		
	aattcaagct tgacgtcagg tggcactttt cggggaaatg tgcgcggaac ccctatttgt	60	
	ttatitttct aaatacattc aaatatgtat ccgtcatga gacaataacc ctgataaatg	120	
50	cttcaataat attgaaaaag gaagagtagt agtattcaac atttccgtgt cgccttatt	180	
	cccttttttg cggcattttg ccttccgtgt ttgtctacc cagaaacgt ggigaaagta	240	
55			

aaagatgctg aagatcagtt ggggtgcacga gtgggttaca tcgaactgga tctcaacagc 300
 5 ggtaagatcc ttgagagttt tcgccccgaa gaacgttttc caatgatgag cactttttaa 360
 gtictgctat gtggcgcggt attatcccggt attgacgccg ggcaagagca actcgggtcgc 420
 cgcatacact attctcagaa tgacttggtt gagtactcac cagtcacaga aaagcatctt 480
 10 acggatggca tgacagtaag agaattatgc agtctgccca taacctagag tgataacact 540
 gcggccaact tactttctgac aacgatcgga ggaccgaagg agctaaccgc ttttttgac 600
 aacatggggg atcatgtaac tcgcttgat cgttgggaac cggagctgaa tgaagccata 660
 15 ccaacgacg agcgtgacac cacgatgcct gtagcaatgg caacaacgtt gcgcaacta 720
 ttaactggcg aactacttac tctagcttcc cggcaacaat taatagactg gatggaggcg 780
 gataaagttg caggaccact tctgcgtcg gcccttccgg ctggttggtt tattgctgat 840
 20 aaatctggag ccggtgagcg tgggtctcgc ggtatcattg cagcactggg gccagatggt 900
 aagccctccc gtatcgtagt tatctacacg acggggagtc aggcaactat ggatgaacga 960
 aatagacaga tcgtgagat aggtgccica ctgattaagc attgtaact gtcagaccaa 1020
 25 gtttactcat atatacttta gattgattta aaacttcatt ttaatttaa aaggatctag 1080
 gtgaagatcc tttttggctc gagtctcatg accaaaatcc cttaacgtga gttttcgttc 1140
 cactgagcgt cagaccccggt agaaaagatc aaaggatctt cttgagatcc ttttttctg 1200
 30 cgcgtaatct gctgcttgca aacaaaaaaa ccaccgtac cagcggtggt ttgtttgccg 1260
 gatcaagagc taccaactct ttttcgaag gtaactggct tcagcagagc gcagatacca 1320
 35 aatactgtcc ttctagtgta gccgtagtta ggccaccact tcaagaactc tgtagaccg 1380
 cctacatacc tcgtctgct aatcctgta ccagtgctg ctgccagtgg cgataagtcg 1440
 tctcttaccg ggttgactc aagacgatag ttaccggata aggcgcagcg gtcgggctga 1500
 40 acgggggggt cgtgcacaca gccagcttg gagcgaacga cctacaccga actgagatac 1560
 ctacagcgtg agcattgaga aagcgcacg cttcccgaa ggagaaaggc ggacaggtat 1620
 ccggtaaagc gcagggtcgg aacaggagag cgcacgagg agcttcagg gggaaacgcc 1680
 45 tggtaicctt atagtcctgt cgggtttcgc cactctgac ttgagcgtc atttttgta 1740
 tgctcgtcag gggggcggag cctatggaaa aacgccagca acgcggcctt ttacggttc 1800
 ctggcctttt gctggccttt tgctcacatg ttcttctctg cgttatcccc tgattctgtg 1860
 50 gataaccgta ttaccgcctt tgagtgaact gataccgctc gccgcagccg aacgaccgag 1920
 cgcagcgagt cagtgaacga ggaagcggaa gagcgcccaa tacgaaacc gcctctcccc 1980

ggcggttggc ctgatcagaa ttcatatgca cgtgttcccg atctagtaac atagatgaca 2040
 5 ccgcgcgcga taatttatcc tagtttgcgc gctatatatt gttttctatc gcgtattaaa 2100
 tgtataattg cgggactcta atcataaaaa cccatctcat aaataacgtc atgcattaca 2160
 tgttaattat tacatgcita acgtaattca acagaaatta tatgataatc atcgcaagac 2220
 10 cggaacagg attcaatctt aagaaacttt attgccaaat gtttgaacga tctgcttcgg 2280
 aggttacctt atctgatttt tgtaaaggtc tgataatggt ccgttgtttt gtaaatcagc 2340
 cagtcgcttg agtaaagaat ccggtctgaa tttctgaagc ctgatgtata gttaatatcc 2400
 15 gcttcacgcc atgttcgtcc gcttttgcgc gggagtttgc cttccctgtt tgagaagatg 2460
 tctccgccga tgcctttccc cggagcgcgc tctgcaaggt tcccttttga tgccaccagc 2520
 ccgagggctt gtgcttctga ttttgaatg taattatcag gtagcttatg atatgtctga 2580
 20 agataatccg caaccccgtc aaacgtgttg ataaccggtc ccatcgcgac ggcttgatgg 2640
 atctcttgcg ggacaccggg atgctaggat gggttatcgt ggccggcgctg cgtgtgtggc 2700
 25 ttttgtagge gccggcgagc gcgggggcaa tgtggcaggt gagtcacggt gcaagcgtgc 2760
 gcaagtgaat gcaacaacca aggacggtca tggcgaaagc acctcacgcg tccaccgtct 2820
 acaggatgta gcagtagcac ggtgaagaa gtgttgctcc gtccattagg tgcattctca 2880
 30 ccgttggcca gaacaggacc gttaacagt taggttgagt gtaggacttt tacgttggtta 2940
 atgtatggca aatagtagta aattttgccc ccattggtct ggctgagata gaacatatc 3000
 tggaaagcct ctagcatatc tttttgaca gctaaacttt gcttcttgcc ttcttggctt 3060
 35 agcaatgacg ttgcccatgt cgtggcaaac atctggtaag gtaactgtat tgcgttgttc 3120
 ccttcaacgg ctcaatcccc acaggccaag ctatcctttc cttggcagta taggctcctt 3180
 gagagattat actaccattt ttaagtgctt ataaagacga tgcctcttaa ccagatcgat 3240
 40 cagaaacaca aagttttagc agcgtaatat cccacacaca tacacacacg aagctatgcc 3300
 tcttcatttt ccgagagatt ctgacagtga ccagaatgtc agaatgccat ttcattggga 3360
 caagtcgac cacaagcttc ttggtggagg tcaagggtgt ctattattat tcgctttcta 3420
 45 ggaaattatt cagaattagt gccttttate ataatcttc tctgagccga tgtggttttg 3480
 gatttcattg ttgggagcta tgcagttgcg gatattctgc tgtggaagaa caggaaactta 3540
 tetgcggggg tcttgcgtgg ggcaacattg atatggttcc tgttcgatgt agtagaatac 3600
 50 aatataattc cgtccttttg ccagattgcc attcttgcca tgcttggat cttcatttgg 3660
 tcaaatgccg caccactctt ggacaggat tagctttatt tctgtggag atggtagaaa 3720

55

actcagctta cagaaatggc atttcacgta gtataacgca agacattagg tactaaaact 3780
 5 caactaactg ttccgaatt tcagggcccc tccaaggatc ccagaaatca tcattctctga 3840
 acatgccttc agagaaatgg cattgaccgt ccattacaaa ctaacgtaca ctgtatctgt 3900
 tctttacgac attgcatgtg gaaaggatct gaagagattt ctcttggtac ataataatct 3960
 10 actcctttgc tacgttaata agagatgtaa aaacatgcaa cagttccagt gccaacattg 4020
 tccaaggatt gtgcaattct ttctggagcg ctaaaatiga ccagattaga cgcacagaa 4080
 tattgaattg cagagttagc caataatcct cataatgtta atgtgctatt gtgtttcact 4140
 15 actcaatata gtcttggaact aacaatcaga ttgtttatga tattaagggtg gttggaictc 4200
 tattggtaatt gtcggcgatt ggaagttctt gcagcttgac aagtctacta tatattggta 4260
 ggtattccag ataaatatta aattttaata aaacaatcac acagaaggat ctgcggccgc 4320
 20 tagcctaggc ccggggccac aaaaatctga gcttaacagc acagttgctc ctctcagagc 4380
 agaatcgggt attcaacacc ctcatatcaa ctactacgtt gtgtataacg gtccacatgc 4440
 cggtatatac gatgactggg gttgtacaaa ggcggaaca aacggcgctc ccggagtgc 4500
 25 acacaagaaa ttgccacta ttacagagcg aagagcagca gctgacgcgt acacaacaag 4560
 tcagcaaaca gacagggtga acttcatccc caaaggagaa gctcaactca agcccaagag 4620
 ctttgetaag gcctaacaa gccaccaaaa gcaaaaagcc cactggetca cgttaggaac 4680
 30 caaaggccc agcagtgatc cagcccaaaa agagatctcc ttgccccgg agattacaat 4740
 ggacgatttc ctctatcttt acgatctagg aaggaagttc gaaggtaag gtgacgacac 4800
 35 tatgttcacc actgataatg agaaggtag cctcttcaat ttcagaaga atgctgaccc 4860
 acagatggtt agagaggcct acgcagcagg tctcatcaag acgatctacc cgagtaacaa 4920
 tctccaggag atcaaatacc ttccaagaa ggttaagat gcagtcaaaa gattcaggac 4980
 40 taattgcac aagaacacag agaagacat atttctcaag atcagaagta ctattccagt 5040
 atggacgatt caaggcttgc ttcataaacc aaggcaagta atagagattg gactctctaa 5100
 aaaggtagtt cctactgaat ctaaggccat gcatggagtc taagattcaa atcgaggatc 5160
 45 taacagaact cgccgtgaag actggcgaac agttcataca gactctttta cgactcaatg 5220
 acaagaagaa aatcttcgtc aacatggtgg agcacgacac tctggtctac tccaaaaatg 5280
 tcaaagatac agtctcagaa gaccaagggt ctattgagac tttcaacaa aggataattt 5340
 50 cgggaacct cctcggttc cattgccag ctatctgtca ctcatcgaa aggacagtag 5400
 aaaaggaagg tggtctctac aatgccatc attgcgataa aggaaggct atcattcaag 5460

55

atgcctctgc cgacagtggc cccaaagatg gaccccccacc cagcaggagc atcgtggaaa 5520
 5 aagaagacgt tccaaccacg tcttcaaagc aagtggattg atgtgacatc tccactgacg 5580
 taaggatga cgcacaatcc cactatcctt cgcaagaccc ttctctata taaggaagtt 5640
 catttcattt ggagaggaca cgctgaaatc accagtctct ctctataaat ctatctctct 5700
 10 ctctataacc atggacccag aacgacgccc ggccgacatc cgcctgcca ccgaggcggg 5760
 catgccggcg gtctgcacca tcgtcaacca ctacatcgag acaagcacgg tcaacttcg 5820
 taccgagccg caggaaccgc aggagtggac ggacgacctc gtccgtctgc gggagcgcta 5880
 15 tccctggctc gtcccgagg tggacggcga ggtccggcg atcgctacg cgggcccctg 5940
 gaaggcacgc aacgcctacg actggacggc cgagtcgacc gtgtacgtct cccccgcc 6000
 ccagcggacg ggactgggct ccacgtctta caccacctg ctgaagtccc tggaggcaca 6060
 20 gggcttcaag agcgtggctc ctgtcatcgg gctgcccaac gacccgagcg tgcgcatgca 6120
 cgaggcgctc ggatatgccc cccgcggcat gctcggggcg gccggcttca agcacgggaa 6180
 ctggcatgac gtgggtttct ggacgtgga cttcagcctg ccgtaccgc cccgtccggt 6240
 25 cctgcccgtc accgagatct gagatcacgc gtcttaggat ccccgatga gctaagctag 6300
 ctatatcatc aatttatgta ttacacataa tatcgactc agtctttcat ctacggcaat 6360
 gtaccagctg atataatcag ttattgaaat atttctgaat ttaaacttgc atcaataaat 6420
 30 ttatgttttt gcttggacta taatacctga cttgttattt tatcaataaa tatttaact 6480
 atatttcttt caagatggga attaacatct acaaatigcc tttcttate gaccatgtac 6540
 35 gtatcgcg 6548

<210> 5

40 <211> 6539

<212> DNA

<213> Escherichia coli LE392

45 <220>

<223> Clone: pTS431

50 <400> 5

aattcaagct tgacgtcagg tggcactttt cgggaaatg tgcgcggaac ccctatttgt 60

55

ttatTTTTct aaatacattc aaatatgtat ccgctcatga gacaataacc ctgataaatg 120
 5 cticcaataat attgaaaaag gaagagtatg agtattcaac atttccgtgt cgcccttatt 180
 cccttttttg cggcattttg ccttccgtgt ttigtccacc cagaaacgct ggtgaaagta 240
 aaagatgctg aagatcagtt ggggtcacga gtgggttaca tcgaactgga tctcaacagc 300
 10 ggtaagatcc ttgagagttt tcgccccgaa gaacgttttc caatgaigag cactttttaa 360
 gttctgctat gtggcgcggt attatcccggt attgacgccg ggcaagagca actcgggtcg 420
 cgcatacact attctcagaa tgacttggtt gagtactcac cagtcacaga aaagcatctt 480
 15 acggatggca tgacagtaag agaattaigc agtgcgccca taaccatgag tgataaacact 540
 gcggccaact tacttctgac aacgatcgga ggaccgaagg agctaaccgc tttttgca 600
 aacatggggg atcatgtaac tcgcttgat cgttgggaac cggagctgaa tgaagccata 660
 20 ccaaacgacg agcgtgacac cacgatgcct gtagcaatgg caacaacggt gcgcaacta 720
 ttaactggcg aactacttac tctagcttcc cggcaacaat taatagactg gatggaggcg 780
 gataaagttg caggaccact tctgcgtcg gcccttccgg ctggctggtt tattgctgat 840
 25 aaatctggag ccggtgagcg tgggtctcgc ggtatcattg cagcactggg gccagatggt 900
 aagccctccc gtatcgtagt tatctacag acggggagtc aggcaactat ggatgaacga 960
 aatagacaga tcgtgagat aggtgcctca ctgattaage attgtaact gtcagaccaa 1020
 30 gtttactcat atatacttta gattgattta aaacttcatt ttaatttaa aaggatctag 1080
 gtgaagatcc ttttggctc gagtctcatg accaaaatcc cttaacgiga gtttctgtc 1140
 cactgagcgt cagaccccggt agaaaagatc aaaggatctt cttgagatcc ttttttctg 1200
 35 cgcgtaatct gctgcttgca aacaaaaaaa ccaccgctac cagcgggtgt ttgtttgccg 1260
 gatcaagagc taccaactct ttttccgaag gtaactggct tcagcagagc gcagatacca 1320
 40 aatactgtcc ttctagtta gccgtagtta ggccaccact tcaagaactc tctagcaccg 1380
 cctacatacc tcgtctgct aatcctgta ccagtggctg ctgccagtgg cgataagtcg 1440
 tgtcttaccg ggttgactc aagacgatag ttaccggata aggcgcagcg gtcgggtga 1500
 45 acgggggggt cgtgcacaca gccagcttg gagcgaacga cctacaccga actgagatac 1560
 ctacagcgtg agcatigaga aagcgccacg ctccccgaag ggagaaaggc ggacaggtat 1620
 ccggtaaagc gcagggtcgg aacaggagag cgcacgaggg agcttcagg gggaaacgcc 1680
 50 tggatatctt atagtctgt cgggttctgc cacctctgac ttgagcgtcg atttttgta 1740
 tgctcgtcag gggggcggag cctatggaaa aacgccagca acgcgccctt ttacggttc 1800

EP 1 020 527 A1

ctggcctttt gctggccttt tgetcacatg ttttttctg cgttatcccc tgattctgtg 1860
5 gataaccgta ttaccgcctt tgagttagct gataccgctc gccgcagccg aacgaccgag 1920
cgcagcgagt cagttagcga ggaagcgaa gagcgccaa tacgcaaacc gcctctcccc 1980
gcgcgttggc ctgatcagaa ttttccccga tctagtaaca tagatgacac cgcgcgcgat 2040
10 aatttatcct agtttgcgcg ctatattttg ttttctatcg cgtattaaat gtataattgc 2100
gggactctaa tcataaaaac ccatctcata aataacgtca tgcatatcat gttaattatt 2160
acatgcttaa cgtaattcaa cagaaattat atgataatca tcgcaagacc ggcaacagga 2220
15 ttcaatctta agaaacttta ttgccaaatg ttgaaacgat ctgcttcgga tctcttagat 2280
tactgatttt tgtaaaggtc tgataatggt ccgttgtttt gtaaatacagc cagtgcgttg 2340
agtaaagaat ccggtctgaa tttctgaagc ctgatgtata gtaatatatc gcttcacgcc 2400
20 atgttcgtcc gcttttggcc gggagttagc ctccctgtt tgagaagatg tctccgccga 2460
tgcttttccc cggagcgacg tctgcaaggt tcccttttga tgccaccag ccgagggtt 2520
gtgcttctga ttttgtaatg taattatcag gtagcttatg atatgtctga agataatccg 2580
25 caaccccgtc aaacgtgttg aataaccggt accatcgca cggcttgatg gatctcttgc 2640
tgacacccgg gatgctagga tgggttatcg tggccggcgt gcgtgtgtgg cttttgtagg 2700
cgccggcgac ggccggggca atgtggcagg tgagtcacgg tgcaagcgtg cgcaagtac 2760
30 tgcaacaacc aaggacggtc atggcgaaag cactcacgc gtccaccgtc tacaggatgt 2820
agcagtagca cggtgaaaga agtgtgtcc cgtccattag gtgcattctc accgttggcc 2880
35 agaacaggac cgttcaacag ttaggttagg tgtaggactt ttacgtggtt aatgtatggc 2940
aaatagtagt aaattttgcc ccattggtc tggctgagat agaacaattt ctggaaagcc 3000
tctagcatat cttttttgac agctaaactt tgcttcttgc cttcttggtc tagcaatgac 3060
40 gttgccatg tctgggcaaa catctggtaa ggtaactgta ttcgtttgtt cccttcaacg 3120
gctcaatccc cacaggccaa gctatccttt ccttggcagt ataggtcctt tgagagatta 3180
tactaccatt ttttaagtct tataaagacg atgctctcta accagatcga tcagaaacac 3240
45 aaagttttag cagcgttaata tcccacacac atacacacac gaagctatgc ctctcattt 3300
tccgagagat tctgacagt accagaatgt cagaatgcca tttcatgggc acaagtcgat 3360
ccacaagctt cttggtagg gtcaagggt gctattatta ttcgtttctt aggaattat 3420
50 tcagaattag tgccctttat cataactct ctctgagccg atgtggtttt ggatttcatt 3480
gttgggagct atgcagttgc ggatattctg ctgtggaaga acaggaaactt atctcgggg 3540

55

gtccttgctg gggcaacatt gatatggtc ctgttcgatg tagtagaata caatataatt 3600
 5 cgcctccttt gccagattgc cattcttggc atgcttgga tcttcatttg gtcaaagcc 3660
 gcaccactct tggacaggta ttagctttat ttcctgtgga gatggiagaa aactcagctt 3720
 acagaaatgg catttcacgt agtataacgc aagacattag gtactaaaac tcaactaact 3780
 10 gtttcgaat ttcagggcc ctccaaggat cccagaaatc atcatctctg aacatgcctt 3840
 cagagaaatg gcattgaccg tccattacaa actaacgtac actgtatctg tttttacga 3900
 cattgcatgt ggaaggatc tgaagagatt tctcctgga cataataatc tactcctttg 3960
 15 ctacgttaat aagagatgta aaaacatgca acagttccag tgccaacatt gtccaaggat 4020
 tgttcaattc tttctggagc gctaaaattg accagattag acgcatcaga atattgaatt 4080
 gcagagttag ccaataatcc tcataatgtt aatgtgctat tgttggtcac tactcaatat 4140
 20 agttctggac taacaatcag attgtttatg atattaaggt ggttggatct ctattggtat 4200
 tgtcggcgat tggaagttct tgcagcttga caagtctact atatattggt aggtattcca 4260
 gataaatatt aaattttaat aaaacaatca cacagaagga tctgcggcgc ctacgcctagg 4320
 25 cccgggccca caaaaatctg agcttaacag cacagttgct cctctcagag cagaatcggg 4380
 tattcaacac cctcatatca actactacgt tgtgtataac ggtccacatg ccggtatata 4440
 cgatgactgg ggttgtaaca aggcggcaac aaacggcggt cccggagttg cacacaagaa 4500
 30 atttgccact attacagagg caagagcagc agctgacgcg tacacaacaa gtcagcaaac 4560
 agacagggtg aacttcatcc ccaaggaga agctcaactc aagcccaaga gctttgctaa 4620
 35 ggccctaaca agcccacaa agcaaaaagc cacttgctc acgctaggaa ccaaaaggcc 4680
 cagcagtgat ccagcccaa aagagatctc ctttgccccg gagattacaa tggacgattt 4740
 cctctatctt tacgatctag gaaggaagt cgaaggtaga ggtgacgaca ctatgttcac 4800
 40 cactgataat gagaaggta gcctcttcaa tttcagaaag aatgctgacc cacagatggt 4860
 tagagaggcc tacgcagcag gtctcatcaa gacgatctac ccgagtaaca atctccagga 4920
 gatcaaatac cttccaaga aggttaaga tgcagtcaaa agattcagga ctaattgcat 4980
 45 caagaacaca gagaagaca tttttctcaa gatcagaagt actattccag tatggacgat 5040
 tcaaggcttg cttcataaac caaggcaagt aatagagatt ggagtctcta aaaaggtagt 5100
 tctactgaa tctaaggcca tgcattgagt ctaagattca aatcgaggat ctaacagaac 5160
 50 tcgccgtgaa gactggcgaa cagttcatc agagtctttt acgactcaat gacaagaaga 5220
 aaatcttcgt caacatggtg gagcagcaca ctctggicta ctccaaaaat gtcaaagata 5280

5 cagtctcaga agaccaaagg gctattgaga cttttcaaca aaggataatt tcgggaaacc 5340
 tcctcggatt ccattgccca gctatctgtc acttcatcga aaggacagta gaaaaggaag 5400
 gtggctccta caaatgccat catcgcgata aaggaaaggc tatcattcaa gatgcctctg 5460
 ccgacagtgg tcccaaagat ggacccccac ccacgaggag catcgtggaa aaagaagacg 5520
 10 ttccaaccac gtcttcaaag caagtggatt gatgtgacat ctccactgac gtaagggatg 5580
 acgcacaatc ccactatcct tcgcaagacc ctctctctat ataaggaagt tcatttcatt 5640
 tggagaggac acgtgaaat caccagtctc tctctataaa tctatctctc tctctataac 5700
 15 catggacca gaacgacgcc cggccgacat ccgccgtgcc accgaggcgg acatgccggc 5760
 ggctgcacc atcgtcaacc actacatcga gacaagcacg gtcaacttcc gtaccgagcc 5820
 gcaggaaccg caggagtggc cggacgacct cgtccgtctg cgggagcgt atccctggct 5880
 20 cgtcgcggag gtggacggcg aggtcgcgg catgcctac gggggcccct ggaaggcacg 5940
 caacgcctac gactggacgg ccgagtcgac cgtgtacgtc tccccccgcc accagcggac 6000
 25 gggactgggc tccacgtct acacccacct gctgaagtcc ctggaggcac agggcttcaa 6060
 gagcgtggtc gctgtcatcg ggctgcccaa cgacccgagc gtgcgcatgc acgaggcgt 6120
 cggatatgcc ccccgcgga tgcctggggc ggccggcttc aagcacggga actggcatga 6180
 30 cgtgggtttc tggcagctgg acttcagcct gccggtaccg ccccgtcgg tctgcccgt 6240
 caccgagatc tgagatcacg cgttctagga tccccgatg agctaagcta gctatatcat 6300
 caatttatgt attacacata atatcgcaat cagtcttica tctacggcaa tgtaccagct 6360
 35 gatataatca gttattgaaa tatttctgaa tttaaacttg catcaataaa tttatgtttt 6420
 tgcttggact ataatacctg acttggtatt ttatcaataa atatttaaac tatatttctt 6480
 40 tcaagatggg aattaacatc tacaattgc cttttcttat cgaccatgta cgtatcgcg 6539

<210> 6

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer 1

<400> 6

5 cgttcggctc gatggtaccg gttatcaaca cgtttga 37

10 <210> 7

<211> 38

<212> DNA

15 <213> Artificial Sequence

<220>

20 <223> Primer 2

25 <400> 7

cctctagatt atctgatttt tgtaaaggtc tgataatg 38

30

Claims

- 35 1. A mutant barnase gene having one or some mutations at least in part of the DNA sequence of barnase gene, wherein said mutant barnase gene is capable, when anther-specifically expressed in a plant, of making said plant substantially male sterile without exerting any substantially disadvantageous effect on the tissues except for the anthers.
- 40 2. The gene as claimed in Claim 1, wherein said mutation is a mutation causing frame shift re-coding.
3. A gene which is derived from a DNA sequence encoding the same amino acid sequence as shown in SEQ ID NO:1 by substitution, deletion, insertion or addition of one to several nucleotides in said DNA sequence, wherein said gene is capable of encoding a protein which makes a plant substantially male sterile when expressed anther-specifically in said plant.
- 45 4. A gene comprising a DNA sequence which is derived from a DNA sequence encoding the same amino acid sequence as shown in SEQ ID NO:1 by substitution, deletion, insertion or addition of one to several nucleotides in the latter DNA sequence, and a promoter located upstream of said DNA sequence for allowing an anther-specific expression, wherein said gene is capable of making a plant substantially male sterile when introduced into the genome of said plant.
- 50 5. A gene represented by SEQ ID NO:3 which encodes a protein capable of making a plant substantially male sterile when expressed anther-specifically in said plant.
- 55 6. A gene comprising the sequence represented by SEQ ID NO:3 and a promoter located upstream of said sequence for allowing an anther-specific expression, wherein said gene is capable of making a plant substantially male sterile when introduced into the genome of said plant.

EP 1 020 527 A1

7. A recombinant vector which contains a gene as claimed in any of Claims 1 to 6 and expresses said gene in a host plant.
8. A method of making a plant male sterile which comprises transforming said plant by a mutant barnase gene as
5 claimed in any of Claims 1 to 6 and allowing said mutant barnase gene to be expressed anther-specifically.
9. The method as claimed in Claim 8, wherein said plant is transformed by said mutant barnase gene by integrating said gene into the genome of said plant.
- 10 10. A transgenic plant wherein a gene as claimed in any of Claims 1 to 6 has been introduced.

15

20

25

30

35

40

45

50

55

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP99/04167

A. CLASSIFICATION OF SUBJECT MATTER

Int.Cl⁶ C12N15/55, C12N9/22, C12N15/82, A01H1/00, A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int.Cl⁶ C12N15/55, C12N9/22, C12N15/82, A01H1/00, A01H5/00

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
WPI/BIOSIS (DIALOG), REGISTRY (STN)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO, 96/26283, A (Plant Genetic Systems N.V.), 29 August, 1996 (29. 08. 96) & EP, 811070, A1 & JP, 11-500617, A	1-10
A	Journal of Immunology Vol. 147 No. 8 (1991) Fettes J.V. et al., "A frameshift mutation at the amino terminus of the nucleoprotein gene does not affect generation of cytotoxic T lymphocyte epitopes" p.2697-2705	1-10

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date
document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"T" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"a" document member of the same patent family

Date of the actual completion of the international search
28 October, 1999 (28. 10. 99)Date of mailing of the international search report
9 November, 1999 (09. 11. 99)Name and mailing address of the ISA/
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

